VARIATIONS ON MURMURATIONS

KIMBALL MARTIN

ABSTRACT. We explore several variations on the recently discovered phenomena of murmurations for elliptic curves and modular forms.

CONTENTS

1.	Introduction	1
2.	Review of murmurations	2
3.	No root numbers	3
4.	Möbius sums	5
5.	Local root numbers	6
6.	Class number sums	8
7.	<i>L</i> -values	9
8.	Representations by quadratic forms	12
References		14

1. INTRODUCTION

Murmurations are a subtle correlation in a sufficiently large family of objects, such as elliptic curves or modular forms, between the root number/rank of the objects and Fourier/Dirichlet coefficients. They were first discovered numerically in the context of elliptic curves by He, Lee, Oliver, and Pozdnyakov [HLOP]. Later, extensive computations of Sutherland with ideas of others elucidated the patterns in murmurations, and indicated such murmurations exist for other families of objects as well, such as modular forms and genus 2 curves – e.g., see slides or data on Sutherland's website.¹ Zubrilina has proved the existence and some properties of murmurations in the context of modular forms of squarefree level [Zub]. Murmurations have also been exhibited in several other settings, such as Dirichlet characters [LOP25], Maass forms [BLLD⁺], and higher rank zeta functions of elliptic curves [SW].

In this note, I will explore some different kinds of variants of murmurations, primarily from an empircal perspective. We will go beyond the paradigm of looking at correlations between root numbers and Fourier/Dirichlet coefficients, and think about the following perspectives:

- (1) expected minus actual solution counts
- (2) averaging arithmetic functions over subsets of integers

Date: May 2, 2025.

¹https://math.mit.edu/~drew/.

(3) traces of "arithmetic" linear operators, and generalizations

Note that a_p 's of elliptic curves are of type (1). Averages of Fourier coefficients of modular forms are traces of Hecke operators, i.e., type (3), which via the trace formula can be expressed as class number sums, i.e., type (2). The "and generalizations" in (3) refers to considering alternate types of trace formulas, such as Kuznetsov or relative trace formulas, as opposed to Eichler–Selberg-type trace formulas.

Acknowledgements. This note is largely based on presentations at the workshops Murmurations in Arithmetic (ICERM, July 6–8, 2023) and Murmurations in Arithmetic Geometry and Related Topics (Simons Center, Nov 11–15, 2024), as well as ensuing discussions with participants. For this I am grateful to the organizers. I also thank Alex Cowan, Thomas Oliver, Andrew Sutherland, and Nina Zubrilina for several helpful discussions and comments.

2. Review of murmurations

Let \mathcal{F} be one of the following two types of families:

- ${\mathcal E}$ elliptic curves up to isogeny, partially ordered by conductor N
- \mathcal{H}_k weight k cuspidal newforms with trivial nebentypus, partially ordered by level N

Fix $\beta > 1$; by default we take $\beta = 2$ following Sutherland. Let $\mathcal{F}^{\pm}(N)$ be the set objects $f \in \mathcal{F}$ with conductor/level N and root number ± 1 . If $\mathcal{F} = \mathcal{E}$, set k = 2. Consider the averages

$$\mathcal{A}_{\mathcal{F}}^{\pm}(p,X) = \mathcal{A}_{\mathcal{F}}^{\pm}(p,X;\beta) = p^{1-\frac{k}{2}} \frac{\sum_{X \le N \le \beta X} \sum_{f \in \mathcal{F}^{\pm}(N)} a_p(f)}{\sum_{X \le N \le \beta X} \# \mathcal{F}^{\pm}(N)}.$$

The $p^{1-\frac{k}{2}}$ is a normalization factor, which is 1 if $\mathcal{F} = \mathcal{E}$ or \mathcal{H}_2 . Here the prime on the sums over N denotes a possible restriction on the N considered — in this section we will take N squarefree and coprime to p, largely as a computational convenience.

For $\mathcal{F} = \mathcal{H}_2$, we plot $\mathcal{A}_{\mathcal{F}}^{\pm}(p; X)$ as a function of p for X = 1000 in Fig. 1 and X = 2000in Fig. 2; the case of root number +1 is plotted in blue and root number -1 in red. Note that these graphs look essentially the same, even though the underlying data comes from completely disjoint sets of newforms! We have chosen the horizontal scale to range up to $p \leq 4X$ in both cases, so one might conjecture that if we continue to make such graphs for larger and larger X, they will tend to the graph of a smooth function in $\frac{p}{X}$, called the *murmuration function*. This limiting property is called *scale invariance* in $\frac{p}{X}$ (or $\frac{p}{N}$).

Indeed this conjecture holds for any even $k \ge 2$ and any $\beta > 1$, as shown by Zubrilina [Zub] with careful estimates of class number sums coming from trace formulas. Zubrilina also describes the resulting murmuration functions. Numerically, the same phenomenon happens without the N squarefree restriction (the coprime to p restriction is asymptotically negligible), but the trace formula is more complicated and the murmuration conjecture has not been proved in this case.

For the case of elliptic curves, again one numerically sees a sort of scale invariance in $\frac{p}{X}$, but there is also noise in the graph that does not seem to disappear, and a more reasonable conjecture is that the averages $\mathcal{A}_{\mathcal{E}}^{\pm}(p;X)$ only tend to a function in $\frac{p}{X}$ after suitable smoothing (e.g., smooth by averaging over nearby p for each X). The other thing to notice about the graphs in Figs. 1 and 2 is that the blue and red (i.e., + and -) graphs are almost reflections of each other. Indeed the limiting murmuration functions for opposite root numbers are precisely negatives of each other, so for asymptotic purposes one may simply consider the difference

$$\mathcal{A}_{\mathcal{F}}^{\Delta}(p,X) = \mathcal{A}_{\mathcal{F}}^{+}(p,X) - \mathcal{A}_{\mathcal{F}}^{-}(p,X).$$

This has the advantage that the inner sum in the numerator for $\mathcal{A}_{\mathcal{F}}^{\Delta}(p, X)$ may now (in the case of $\mathcal{F} = \mathcal{H}_k$) be expressed as the trace

$$\operatorname{tr}_{\mathcal{F}(N)} WT_p := \sum_{f \in \mathcal{F}(N)} w_f a_p(f) = \sum_{f \in \mathcal{F}^+(N)} a_p(f) - \sum_{f \in \mathcal{F}^-(N)} a_p(f),$$

where W is the Fricke involution on the newspace $S_k^{\text{new}}(N)$ and w_f is the root number of f.² Hence one can analyze such averages now with the trace formula. We note that in the simple case of k = 2, N > 1 squarefree coprime to p, the trace formula reads

(2.1)
$$\operatorname{tr}_{S_2^{\operatorname{new}}(N)} WT_p = \frac{1}{2} \sum_{s^2 \le \frac{4p}{N}} H(s^2 N^2 - 4Np) - (p+1),$$

where $s \in \mathbb{Z}$, and H is the Hurwitz class number.

Here are some additional remarks: (1) This correlation between a_p 's and root numbers is something one only sees averaging over large families, and is quantitatively quite small — after the normalization by $p^{1-\frac{k}{2}}$, each individual Fourier coefficient has size on the order of \sqrt{p} . (2) The Birch and Swinnerton-Dyer conjecture asserts a subtle correlation between sizes of a_p 's and ranks of elliptic curves. (3) For small p, a correlation between a_p and the root number of newforms was already observed in [MP22]; the "limiting case" of p = 1 corresponds to the bias of root numbers toward +1 from [Mar18, Mar23]. (4) One can more generally consider correlation of a_n 's, but for simplicity we stick to a_p 's this restriction makes patterns more apparent with less data.

3. No root numbers

One of the first variations that might come to mind is simply to consider averages without fixing root numbers, which is essentially the average of $\mathcal{A}_{\mathcal{F}}^+(p, X)$ and $\mathcal{A}_{\mathcal{F}}^-(p, X)$. As remarked above, they will tend to 0, but one can still look for structure in the behavior as $X \to \infty$, which can be thought of as a second-order term in comparing the convergence to murmuration functions for root number +1 versus root number -1.

In [Mar], I conjectured the existence such murmurations without root numbers for $\mathcal{F} = \mathcal{H}_k$, but not $\mathcal{F} = \mathcal{E}$. Specifically, consider the weighted averages

$$\mathcal{A}_{\mathcal{F}}(p,X) = p^{1-\frac{k}{2}} \frac{\sum_{X \le N \le \beta X}^{\prime} \sqrt{N} \sum_{f \in \mathcal{F}(N)} a_p(f)}{\sum_{X \le N \le \beta X}^{\prime} \# \mathcal{F}(N)}$$

where $\mathcal{F}(N)$ is the set of $f \in \mathcal{F}$ of level N. Again, for simplicity, the prime on the outer sums means in our calculations we restrict to squarefree N coprime to p, and we take $\beta = 2$. Here we have inserted a scaling factor of \sqrt{N} in the numerator to prevent the averages from going to 0.

²Technically one usually considers the averages of $w_f a_p(f)$ over \mathcal{F} , rather than averaging separately over each root number, but this is approximately the same as $\frac{1}{2}\mathcal{A}_F^{\Delta}(p,X)$.



See Figs. 3 and 4 for graphs of $\mathcal{A}_{\mathcal{F}}(p, X)$ with $\mathcal{F} = \mathcal{H}_2, p \leq 4X$, where X = 2000 and X = 4000, respectively. Note as in the original case of murmurations, there appears to be scale invariance in $\frac{p}{N}$, but there is more noise in the graphs. Graphs without the \sqrt{N} scaling look similar in shape, but have vertical range that tends to 0 as $X \to \infty$. Graphs in higher weight look fairly similar, and for $\mathcal{F} = \mathcal{H}_k$ I conjectured that such graphs tend to a limiting murmuration function after appropriate smoothing.

On the other hand, for elliptic curves, previous calculations of Sutherland indicate no such murmurations if one omits root numbers. See Figs. 5 and 6 for the elliptic curve analogues of Figs. 3 and 4, but with ranges restricted to $p \leq 2X$.

To indicate the difference with the usual murmurations from Section 2, we write down the trace formula for the inner sum $\operatorname{tr}_{\mathcal{F}(N)} T_p = \sum_{f \in \mathcal{F}(N)} a_p(f)$ in the simple case that



 $\mathcal{F} = \mathcal{H}_2$ and N > 1 is squarefree and coprime to p:

(3.1)
$$\operatorname{tr}_{S_2^{\operatorname{new}}(N)} T_p = -\frac{1}{2} \sum_{s^2 \le 4p} \xi_{s^2 - 4p}(N) H(s^2 - 4p) + \mu(N)(p+1),$$

where ξ_{Δ} is a certain multiplicative function defined in [Mar], and μ is the Möbius function. (If q is a prime such that $q^2 \nmid \Delta$, then $\xi_{\Delta}(q) = \left(\frac{\Delta}{q}\right) - 1$.)

A significance difference is that in (3.1) the number of class number terms grows with p, so is unbounded in a limit $p, N \to \infty$, whereas as in (2.1) the number of terms is bounded in $\frac{p}{N}$. This makes a theoretical analysis more challenging.

4. Möbius sums

To try to analyze the averages of a_p 's without root numbers from Section 3 in the case of weight 2 modular forms, one might first try to analyze the contribution of the $\mu(N)(p+1)$ term in (3.1). We remark that this term comes from removing the Eisenstein contribution from the trace of T_p on $M_k(N)$. Specifically, the contribution to $\mathcal{A}_{\mathcal{H}_2}(p, X)$ is

(4.1)
$$\mathcal{A}_{\mathcal{H}_2}^{\mu}(p,X) = (p+1) \frac{\sum_{X \le N \le \beta X}' \sqrt{N} \mu(N)}{\sum_{X \le N \le \beta X}' \dim S_2^{\text{new}}(N)}.$$

For N squarefree, dim $S_2^{\text{new}}(N) = \frac{\varphi(N)}{12} + O(\log N)$, the approximate growth is

(4.2)
$$\mathcal{A}^{\mu}_{\mathcal{H}_2}(p,X) \approx \frac{p}{X^2} \sum_{X \le N \le \beta X} \sqrt{N} \mu(N).$$

(Here \approx means asymptotic up to a scalar.)

The right hand side is a modified version of Mertens function $M(X) = \sum_{1 \le N \le X} \mu(N)$ (or rather a weighted analogue of $M(\beta X) - M(X)$). It is conjectured that $|M(X)/\sqrt{X}|$ is unbounded. (This is subtle—recall that the Riemann hypothesis is equivalent to M(X) =



 $O(X^{\frac{1}{2}+\epsilon})$.) Similarly, one would expect that the sum $\sum' \sqrt{N\mu}(N)$ in the right hand side of (4.2) should not be O(X), and which would mean $\mathcal{A}^{\mu}_{\mathcal{H}_2}(p, X)$ is not bounded as $p, X \to \infty$ such that $\frac{p}{X}$ tends to a non-zero limit.

On the other hand, we just conjectured that $\mathcal{A}_{\mathcal{F}}(p, X)$ is bounded! How can these be compatible? Admittedly, the asymptotics of Mertens-like functions are quite subtle, so perhaps more care should be taken in the above approximation. Still, we suggest that the erratic (in X) behavior of the Möbius contribution $\mathcal{A}^{\mu}_{\mathcal{H}_2}(p, X)$ gets canceled out with the class number sums. That is, we propose there is an almost magical interaction between the $\mu(N)$ term and the class number sum in tr T_p in (3.1). Note that in tr T_p there are roughly \sqrt{p} class numbers appearing, each of order approximately \sqrt{p} , and the $\mu(N)$ term is of size p + 1.

As evidence for this interaction, we decompose the murmuration graphs in Figs. 3 and 4 into the class number contribution (in blue) and the Möbius contribution $\mathcal{A}^{\mu}_{\mathcal{H}_2}(p, X)$ (in red) in Figs. 7 and 8. Namely, summing the red and blue graphs in the latter figures gives the scale-invariant murmration graphs in the former figures. However, neither the class number nor the Möbius contributions individually appear to be scale invariant, as the slope of the Möbius contribution changes with X. See Fig. 9 for a plot of the slopes of the Möbius contribution for $1000 \leq X \leq 4000$.

Lastly, we remark upon the jaggedness of the red lines in these figures: note that the expression for the Möbius contribution in (4.1) is not exactly linear in p due to the fact that we restricted our sums over N to gcd(N, p) = 1.

5. Local root numbers

Both the usual murmurations with respect to root numbers in Section 2 and without root number in Section 3 fall into a more general framework given in [Mar]. Namely, the averages considered in those two situations can be seen as weighted averages of quantities tr WT_p and tr $T_p = \text{tr } W_1T_p$, respectively.

More generally one can consider averages of $\operatorname{tr}_{S_k^{\operatorname{new}}(N)} W_M T_p$'s, where W_M is a suitably chosen Atkin–Lehner operator on $S_k^{\operatorname{new}}(N)$. (In the averages, M will potentially vary with



N; in our notation, the Fricke involution is $W = (-1)^{k/2} W_N$.) Conceptually, one can think of this variant as studying the correlation of Fourier coefficients with (the product of) local root numbers at some subset S of places. Looking at the global root number is the case where S is all places of \mathbb{Q} , and no root number is the case where S is the empty set.

We refer to [Mar] for the full context and details, along with some theoretical results. Here we just illustrate some examples in the simple case of levels of the form N = pq, where p < q are distinct primes. We propose that a good way to study murmurations with respect to local root numbers is to look at murmuration graphs for each Atkin–Lehner eigenspace, analogous to our original murmuration pictures in Figs. 1 and 2 where we plotted graphs for each global root number separately.

For a newform f of level N = pq, the global root number $w_f = \prod_v w_{f,v}$ where $w_{f,v}$ is the local root number at v. Here $w_{f,\infty} = (-1)^{k/2}$, $w_{f,p}, w_{f,q} \in \{\pm 1\}$, and $w_{f,v} = +1$ for all other v. There are 4 Atkin–Lehner eigenspaces, which we denote by the 4 sign patterns ++, +- and -+, --. E.g., +- will refer to the subspace of $S_k^{\text{new}}(N)$ generate by newforms with $w_{f,p} = +1$ and $w_{f,q} = -1$.

We present two types of examples: (i) we fix p and vary q, and (ii) we vary both p and q. See Figs. 10 and 11 for a graphs of averages of a_{ℓ} 's over each Atkin–Lehner eigenspace in $S_2^{\text{new}}(pq)$ and $S_4^{\text{new}}(pq)$. Here p = 2 is fixed and 3000 < q < 6000. See Fig. 12 for $S_4^{\text{new}}(pq)$ where p < q and 6000 < pq < 12000.

In all of these plots, the blue and green dots correspond to signs ++ and -- and red and orange dots to signs +- and -+, respectively. Adding all 4 colors corresponds to looking at murmurations with no root number. Adding just the blue and green (resp., red and orange) graphs corresponds to looking at murmurations for global root number +1 (resp., -1) when k = 4, and -1 (resp. +1) when k = 2.

We remark that if one goes beyond squarefree levels, one can also look at something more refined than just the local root number at ramified places — one can also look at the local inertial type of the representation. Trace formulas that will allow us to do such calculations have been worked out in some cases recently in [Kni].



6. Class number sums

Zubrilina [Zub] proved murmurations for modular forms by deriving asymptotics for short class number sums of the form

$$\sum_{X \le N \le Y}' H(s^2 N^2 - 4Np).$$

This is a sum over suitable levels for a given s-term in (2.1). One can view this as a sum of class numbers $H(f_s(N; p))$ where f_s is a polynomial, and N runs over a restricted range. The existence of murmurations amounts to suitably normalized sums of class numbers having a limit as $p, N \to \infty$ such that $\frac{p}{N} \to x$ for any x > 0.

Numerically we have observed one has similar behavior if one slightly varies the polynomial $f_s(N; p)$, e.g., by varying scalars or adding or subtracting smaller order terms. We do not have a comprehensive philosophy of how such class number sums should behave, or what polynomials are interesting, so we simply present a few numerical examples. We begin by restricting to quadratic polynomials $f_s(N; p)$.

As a baseline, in Fig. 13, we present a graph of

$$A_0(p,X) := \frac{\sum_{X \le N \le 2X} \sum_{s=0}^1 H(s^2 N^2 - 4Np)}{\sum_{X \le N \le 2X} N}$$

for X = 1000 and $p \leq X$. Here \sum' denotes a restriction to squarefree N. This is essentially the s = 0 and s = 1 term contributions to murmurations for $\mathcal{F} = \mathcal{H}_2$ from (2.1). We chose to include both s = 0 and s = 1 terms so that one can see some oscillation — the terms for larger s do not contribute in the range $p \leq X$.

In Fig. 14, we simply consider $f_s(N;p) = s^2p^2 - 4Np$ instead of $s^2N^2 - 4Np$. Here larger s terms come into play in the range $p \leq X$, and we plot

$$A_1(p,X) := \frac{\sum_{X \le N \le 2X} \sum_{s=0}^4 H(s^2 p^2 - 4Np)}{\sum_{X \le N \le 2X} N}$$

for X = 1000. Again there is some oscillatory behavior.



In Fig. 15, we plot

$$A_2(p,X) := \frac{\sum_{X \le N \le 2X} \sum_{s=0}^4 H(s^2 + 1 - 3Np)}{\sum_{X \le N \le 2X} N}$$

for X = 1000. Here each class number discriminant is about size -3Np, and it seems there is not enough variation in adding $s^2 + 1$ to cause oscillation.

In all of the above examples, while we only included graphs for X = 1000, graphs look similar for other values of X, i.e., further calculations indicate scale invariance in $\frac{p}{X}$, i.e., there is a limiting graph as $X \to \infty$.

Now we come to an example of a cubic polynomial $f_s(N; p)$. We plot

$$A_3(p,X) := \frac{\sum_{X \le N \le 2X} \sum_{s=0}^{1} H(sN^3 - 4N^2p)}{\sum_{X \le N \le 2X} N^{3/2}}$$

for X = 1000 in Fig. 16. Note that we changed the normalization factor in the denominator to account for the class numbers now being of size approximately $N^{3/2}$ (assuming $p \approx X$). Again the graphs look roughly similar as $X \to \infty$, but there does not appear to be convergence to an actual function, because we are averaging too sparse a set of class numbers.

But we can modify the last example by averaging over more class numbers. For instance, instead of average over $\approx X$ class numbers for discriminants of size $\approx X^3$, we consider the following average over $\approx X^{3/2}$ such class numbers,

$$A_4(p,X) := \frac{\sum_{X \le N \le 2X} \sum_{t \le \sqrt{X}} \sum_{s=0}^1 H(s^2 N^3 - tN - 4N^2 p)}{\sum_{X \le N \le 2X} \sum_{t \le \sqrt{X}} N^{3/2}}$$

This average seems to converge to a function in $\frac{p}{X}$ — see Fig. 17 for a plot of $A_4(p, X)$ with X = 500.

7. L-VALUES

Several of the murmurations we have considered are just averages of the geometric side of a trace formula. There are other kinds of trace formulas one can consider: the Kunzetsov trace formula, the Petersson trace formula, and more generally relative trace



formulas. The Kuznetsov trace formula will give sums of Fourier coefficients weighted by inverse Petersson norms. We will not address that, but just discuss one different relative trace formula.

One of Jacquet's first relative trace formulas relates toric periods on quaternion algebras to twisted central values of L-functions, reproving a result of Waldspurger. Evaluating the geometric side for the quaternion algebra gives an exact formula for average L-values

$$\Lambda(N,k,D,n) = \sum_{f \in \mathcal{H}_k(N)} \frac{L(1/2,f)L(1/2,f \otimes \chi_{-D})}{(f,f)} a_n(f),$$

where $\chi_{-D} = \left(\frac{-D}{\cdot}\right)$, for some fixed fundamental discriminant -D < 0. This was carried out in [FW09, Mar22] under some hypotheses on N and D.

We just indicate what the geometric side looks like in the following simple setting, which was originally considered in [MR12] (using the Gross–Zagier formula, rather than Jacquet's relative trace formula). Suppose $0 < D \equiv 3 \mod 4$, and $N \neq p$ is a prime inert in $K = \mathbb{Q}(\sqrt{-D})$. Let $h_D = h_K$ and $u_D = [\mathcal{O}_K^{\times} : \mathbb{Z}^{\times}]$. Also assume k = 2 for simplicity. Then

(7.1)
$$\frac{\sqrt{D}u_D^2}{2\pi}\Lambda(N,k,D,n) = \frac{12h_D^2}{N-1}\sigma_N(n) + u_D r(nD)h_D + u_D^2 \sum_{m=1}^{\lfloor nD/N \rfloor} \Phi(m,N),$$

where $\sigma_N(n)$ is the sum of divisors $d \mid n$ such that (d, N) = 1, r(nD) is the number of ideals of norm nD in \mathcal{O}_K and

$$\Phi(m, N) = d((m, D))r(m)r(pD - mN).$$

(Here $d = \sigma_0$ is the number-of-divisors function.)

There are a couple of ways one could investigate murmuration analogues for these quantities. For the standard trace formula for tr T_p or tr $W_N T_p$ on $S_k^{\text{new}}(N)$, there are 3 parameters one can vary: N, k and p, and we varied p and N. (One can also consider varying k—see [BBLLD].) Here there are 4 parameters: N, k, D, n. We choose to keep k



fixed and average over N as in the original murmurations setting, and vary either D or n = p.

First consider varying D. This amounts to considering murmurations for squares of Fourier coefficients of half-integral weight forms (see [Wal81]). In Figs. 18 and 19, we plot averages

$$A_{L,D}(D,X) = \frac{\sum_{X \le N \le 2X}' \Lambda(N,2,D,1)}{\sum_{X \le N \le 2X}' 1} = \frac{\sum_{X \le N \le 2X}' \sum_{f \in \mathcal{H}_2(N)} \frac{L(1/2,f)L(1/2,f \otimes \chi_{-D})}{(f,f)}}{\sum_{X \le N \le 2X}' 1}$$

for X = 4000, 8000, where $-D \equiv 1 \mod 4$ is a negative prime fundamental discriminant. Here the prime in the sum over N refers to a restriction to primes inert in $K = \mathbb{Q}(\sqrt{-D})$ so that (7.1) is valid. We take $3 < D \leq X$ along the horizontal axis, and color the $-D \equiv 1 \mod 8$ points blue and the $-D \equiv 5 \mod 8$ points red.

We tried a few different normalizations/weightings of these averages, and while the graphs look roughly similar as X increases, we did not find a simple one which is scale invariant in $\frac{D}{X}$, or is clearly indicative of murmurations. Note that we divide by the number of levels in the average, rather than the number of newforms, to prevent the graphs from shrinking in scale as X grows. We also remark that such averages only pick up forms with root number +1 because of the L(1/2, f) factor.

Finally, we consider the case of fixing D and varying n = p. Here we consider averaging the scaled quantity in (7.1). Let

$$A_{L,p}(p,X) = \frac{\sum_{X \le N \le 2X}' \Lambda(N,2,3,p)}{\sum_{X \le N \le 2X}' 1} = \frac{\sum_{X \le N \le 2X}' \sum_{f \in \mathcal{H}_k(N)} \frac{L(1/2,f)L(1/2,f \otimes \chi_{-3})}{(f,f)} a_p}{\sum_{X \le N \le 2X}' 1}.$$

In Figs. 20 and 21, we plot $A_{L,p}(p, X)$ for $p \leq X$ $(p \neq 3)$ and X = 2000, 4000, coloring the values blue or red according to $\chi_{-3}(p) = +1$ or -1.

These graphs are apparently scale invariant in $\frac{p}{X}$, and we expect the blue and red graphs to tend to a limiting function after appropriate smoothing. The first linear part of the graph corresponds to the so-called *stable range* $p < \frac{N}{D}$ where the sum on the



right hand side of (7.1) vanishes (in this range, it is not hard to prove the limit exists). This is somewhat analogous to murmurations in Section 2, where the first part of the murmuration graph is simply given by a square root function (or a linear function if one graphs in terms of $\sqrt{\frac{p}{X}}$, which turns out to be nicer), and more trace formula terms contribute the further to the right one goes on the graph.

The reason to separate the cases of p inert or split in $K = \mathbb{Q}(\sqrt{-D})$ it because that affects the second term on the right hand side of (7.1). We also note that $\sum' 1$ is the correct scaling in the denominator to get scale invariance in $\frac{p}{X}$ because the main term (the first term on the right) of (7.1) is of size $\approx \frac{p}{N}$.

8. Representations by quadratic forms

For an elliptic curve E at a good prime p, $a_p(E) = p + 1 - \#E(\mathbb{F}_p)$, which is the deviation of expected minus actual number of solutions mod p. Hence murmurations for elliptic curves indicate how the error in first-order point count estimates mod p is correlated with the conductor.

One other situation in number theory where the error of first-order solution count estimates is well studied is representations by quadratic forms. In fact, both traces of T_p 's and the averages $A_{L,p}$ of *L*-value weighted a_p 's can be interpreted in terms of Brandt matrix entries, which can be expressed as representations numbers of quadratic forms (e.g., see [Mar22, Section 3.3]).

The most classical case is that of representations of integers by binary quadratic forms, which includes the Gauss circle problem. Let -D < 0 be a discriminant, and $r_D(n)$ be the total number of ways to represent n by a positive definite reduced binary quadratic form of discriminant -D. Then

$$r_D(n) = 2u_D \sum_{d|n} \chi_{-D}(n).$$

In particular, for n = p the $\chi_{-D}(p)$ measures the error of the first-order approximation for the number of ways to represent p by a reduced form of discriminant -D. Thus one



analogue of the murmuration averages for elliptic curves in the setting of positive definite binary quadratic forms is the quantity

$$A_{\mathrm{BQF}}^{\mathcal{D}}(p,X) = \frac{1}{\sqrt{X}} \sum_{\substack{X \le D \le 2X \\ -D \in \mathcal{D}}}' \chi_{-D}(p),$$

where \mathcal{D} is a chosen class of fundamental discriminants (see below) and the sum is taken over D coprime to p. For a given p, the sum on the right should be roughly of size \sqrt{X} , which is why we normalize by $\frac{1}{\sqrt{X}}$.

In Figs. 22 and 23 we plot the quantities $A_{BQF}^{\mathcal{D}}(p, X)$ for $p \leq 4X$, where \mathcal{D} is either the set of fundamental discriminants $-D \equiv 1 \mod 8$ (in blue) or $-D \equiv 5 \mod 8$ (in red).

The first plot is with X = 5000 and the second is with X = 10000. See Figs. 24 and 25 for analogous plots where \mathcal{D} is set of fundamental discriminants of the form -4d with either $d \equiv 1 \mod 4$ (in blue) or $d \equiv 3 \mod 4$ (in red). While there is a lot of noise in these graphs, they appear to be roughly scale invariant, which indicates the existence of murmurations in this setting.

Indeed, murmurations for quadratic (and general) Dirichlet characters were already established in [LOP25] under GRH. They smooth out the noisiness by averaging over nearby primes. Very recently, Cowan [Cow] obtained unconditional murmuration results, which also apply to the setting of quadratic Dirichlet characters.

References

- [BBLLD] Jonathan Bober, Andrew R. Booker, Min Lee, and David Lowry-Duda, Murmurations of modular forms in the weight aspect, available at https://arxiv.org/abs/2310.07746. ↑7
- [BLLD⁺] Andrew R. Booker, Min Lee, David Lowry-Duda, Andrei Seymour-Howell, and Nina Zubrilina, Murmurations of Maass forms, available at https://arxiv.org/abs/2409.00765. ¹
 - [Cow] Alex Cowan, On the mean value of GL₁ and GL₂ L-functions, with applications to murmurations, available at https://arxiv.org/abs/2504.09944. ↑8
- [FW09] Brooke Feigon and David Whitehouse, Averages of central L-values of Hilbert modular forms with an application to subconvexity, Duke Math. J. 149 (2009), no. 2, 347–410, DOI 10.1215/00127094-2009-041. MR2541706 ↑7
- [HLOP] Yang-Hui He, Kyu-Hwan Lee, Thomas Oliver, and Alexey Pozdnyakov, Murmurations of elliptic curves, available at https://arxiv.org/abs/2204.10140. Experimental Math, to appear. ↑1
- [Kni] Andrew Knightly, Counting locally supercuspidal newforms, available at https://arxiv.org/ abs/2310.17047. ↑5
- [LOP25] Kyu-Hwan Lee, Thomas Oliver, and Alexey Pozdnyakov, Murmurations of Dirichlet Characters, Int. Math. Res. Not. IMRN 1 (2025), rnae277. [↑]1, 8
- [Mar18] Kimball Martin, Refined dimensions of cusp forms, and equidistribution and bias of signs, J. Number Theory 188 (2018), 1–17. [↑]2
- [Mar22] _____, Exact double averages of twisted L-values, Math. Z. **302** (2022), no. 3, 1821–1854, DOI 10.1007/s00209-022-03088-3. MR4492517 ↑7, 8
- $[Mar23] ____, Root number bias for newforms, Proc. Amer. Math. Soc. 151 (2023), no. 9, 3721–3736.$ $\uparrow 2$
- [Mar] _____, Distribution of local signs of modular forms and murmurations of Fourier coefficients, available at https://arxiv.org/abs/2409.02338. ↑3, 3, 5
- [MP22] Kimball Martin and Thomas Pharis, Rank bias for elliptic curves mod p, Involve 15 (2022), no. 4, 709–726. [↑]2
- [MR12] Philippe Michel and Dinakar Ramakrishnan, Consequences of the Gross-Zagier formulae: stability of average L-values, subconvexity, and non-vanishing mod p, Number theory, analysis and geometry, Springer, New York, 2012, pp. 437–459. ↑7
- [SW] Zhan Shi and Lin Weng, Murmurations and Sato-Tate Conjectures for High Rank Zetas of Elliptic Curves, available at https://arxiv.org/abs/2410.04952. ↑1
- [Wal81] J.-L. Waldspurger, Sur les coefficients de Fourier des formes modulaires de poids demi-entier,
 J. Math. Pures Appl. (9) 60 (1981), no. 4, 375–484 (French). MR0646366 ↑7
 - [Zub] Nina Zubrilina, Murmurations, available at https://arxiv.org/abs/2310.07681. ¹, 2, 6

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019 USA *Email address:* kimball.martin@ou.edu

DEPARTMENT OF MATHEMATICS · INTERNATIONAL RESEARCH AND EDUCATION CENTER, GRADUATE SCHOOL OF SCIENCE, OSAKA METROPOLITAN UNIVERSITY, OSAKA 558-8585, JAPAN

Email address: kimball@omu.ac.jp